

Cinqair (Reslizumab)

Infusion Order Form – Page 1 of 1



Preferred Clinic Location

- ☐ **Harbor Health Park Bend Clinic**
2200 Park Bend Drive, Bldg 2, Suite 300, Austin, TX 78758
P: (855) 481-8375
F: (512) 233-2288

Patient Information

Referral Status: ☐ New Referral ☐ Updated Order ☐ Order Renewal

Date: _____ Patient Name: _____ DOB: _____
ICD-10 code (required): _____ ICD-10 description: _____
☐ NKDA Allergies: _____ Weight (lbs/kg): _____ Height: _____
Patient Status: ☐ New to Therapy ☐ Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

Provider Information

Referral Coordinator Name: _____ Referral Coordinator Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip Code: _____

Supporting Documents/Information (Please provide all of the following)

- Patient insurance information
- Patient medication list
- Supporting clinical notes (H&P) to support primary diagnosis:
 - Lab results showing elevated eosinophil levels
 - List of current medications treating severe asthma
 - Patient has severe asthma with an eosinophilic phenotype and is 18 years of age or older
 - FEV1 test results

Orders

NURSING

- ☒ Nursing care per [Harbor Health Nursing Procedures](#) and [Harbor Health Adverse Reaction Management Protocol](#).

PRE-MEDICATION

- ☐ acetaminophen (Tylenol) ☐ 500mg / ☐ 650mg / ☐ 1000mg PO
☐ cetirizine (Zyrtec) 10mg PO
☐ loratadine (Claritin) 10mg PO
☐ diphenhydramine (Benadryl) ☐ 25mg / ☐ 50mg ☐ PO / ☐ IV
☐ methylprednisolone (Solu-Medrol) ☐ 40mg / ☐ 125mg IV
☐ hydrocortisone (Solu-Cortef) ☐ 100mg IV
☐ Other: _____

Dose: _____ Route: _____
Frequency: _____

THERAPY

- ☒ **Reslizumab** (Cinqair) in 50ml 0.9% sodium chloride intravenous infusion over 25-50 minutes using a 0.2 micron in-line or add-on filter
- Dose: ☐ 3mg/kg
 - ☐ Round up to nearest whole vial
 - ☐ Give exact dose
 - Route: intravenous
 - Frequency: ☐ every 4 weeks ☐ other _____
 - ☒ Flush with 0.9% sodium chloride at infusion completion
- ☒ Patient is required to stay for 30-minute observation

**Consider administering premedication for prophylaxis against infusion reactions and hypersensitivity reactions.*

Provider Name (Print): _____

Provider Signature: _____

Date: _____